

1999

Milkfish breeding and hatchery technology at SEAFDEC/AQD

Aquaculture Department, Southeast Asian Fisheries Development Center

<http://hdl.handle.net/10862/614>

Downloaded from <http://repository.seafdec.org.ph>, SEAFDEC/AQD's Institutional Repository

Milkfish Breeding and Hatchery Technology at SEAFDEC/AQD



AQUACULTURE DEPARTMENT

Southeast Asian Fisheries Development Center

Tigbauan, Iloilo

AQD milkfish technology

supported by more than two decades of research

In support of a sustainable milkfish aquaculture, research on captive breeding and mass fry production were undertaken for the past two decades at SEAFDEC/AQD. These technologies, now being adopted by the private sector, include:

BROODSTOCK MANAGEMENT

Although a scientific breakthrough in 1977, the induced spawning of wild-caught sabalo (adult milkfish) had a low success rate and leads to mortalities. Thus, milkfish were reared to adults in floating net cages (10 m diameter by 3 m deep) and concrete tanks (10 x 10 x 2 m or 10 x 25 x 2 m deep) supplied with flow-through filtered seawater. Sexual maturation began as the broodstock reached 5 years of age (body weight 3 kgs and above). Since then, natural spawning has been observed annually in floating net cages.

Eggs are immediately collected to prevent cannibalism by the broodstock using a manually operated sweeper-type egg collector. Fine mesh knots attached to an airlift system made of 4" diameter PVC pipe collect the eggs in concrete tanks. Egg production of up to 2.5 million/female/season has been attained. Viable eggs are about 80% per spawning, of which 80% hatch. Normal hatchings (straight and without deformities) are usually 80% per spawn.

BROODSTOCK DIET

A diet formulated by SEAFDEC/AQD to support sexual maturation and production of high quality eggs of milkfish is used to feed the broodstock. The diet contains 36% protein (from imported animal and terrestrial proteins), 6% lipid and vitamin mix.

COMMERCIAL FRY PRODUCTION

The water and feeding management schemes for milkfish larviculture have been developed to ensure mass fry production. Hatchery operations utilize either an intensive (high stocking density, high volume tanks, daily feeding and water change) or a semi-intensive (low stocking density, high volume tanks, minimal water change, feeding with mixed diet) system, with an average survival rate of 30% (from stocked newly-hatched larvae). The technology has been used by the private sector to utilize abandoned shrimp hatcheries.



LIVE TRANSPORT

A technique that combines utilizing pre-transport starvation anesthetic at capture, chilled transport water, and a sedating dose of anesthesia effectively transports live broodstock. It was tried on 4-13 year old broodstock placed in oxygenated transport plastic bags (2 m long x 0.5 m wide) containing 40-l chilled (20-22°C) seawater and 5 ml 2-phenoxyethanol (anesthetic). The transport bags were then placed in styro-foam boxes. Travel time was 6-7 hours, recovery was fast, and no mortality was recorded. Eggs (embryonic stage) and newly-hatched larvae were also transported in oxygenated plastic bags containing 12-l seawater and supported by straw bags. Optimum density, temperature, and salinity was 100,000-120,000 eggs or larvae per bag, 28-30 °C and 32-34 ppt.

LARVAL DIET

To minimize the use of mass-cultured rotifers, a formulated larval diet containing adequate nutrition (highly unsaturated fatty acids and vitamin mix) was found to be an effective supplement for rotifers and alternative for the expensive brine shrimp (*Artemia*) nauplii for milkfish larviculture.

PARTIAL LIST OF SEAFDEC/AQD RESEARCH PUBLICATIONS ON MILKFISH

- Agbayani RF, Baliao DD, Franco NM, Ticar RB, Guanzon NG Jr. 1989. An economic analysis of the modular pond system of milkfish production in the Philippines. *AQUACULTURE* 83: 249-259
- Agbayani RF, Lopez NA, Tumaliuan RE, Benjamin GD. 1991 Economic analysis of an integral milkfish broodstock and hatchery operation as a public enterprise. *AQUACULTURE*. 99: 235-248
- Alava VR. 1998. Effect of salinity, dietary lipid source and level of growth of milkfish (*Chanos chanos*) fry. *AQUACULTURE* 167: 229-236
- Alava VR, Kanazawa A. 1996. Effect of dietary fatty acid on growth of milkfish *Chanos chanos* fry in brackishwater. *AQUACULTURE* 144 (4): 363-369
- Almendras JME. 1987. Acute nitrite toxicity and methemoglobinemia in juvenile milkfish (*Chanos chanos* Forsskal). *AQUACULTURE* 61 (1): 33-40
- Bagarinao T. 1994. Systematics, distribution, genetics and life history of milkfish, *Chanos chanos*. *ENVIRON. BIOL. FISHES* 39 (1) 23-41
- Bagarinao T. 1986. Yolk resorption, onset of feeding and survival potential of larvae of three tropical marine fish species reared in the hatchery. *MAR. BIOL.* (91):449-459
- Bagarinao TU, Kumagai S. 1981. Studies on the habitat and food of juvenile milkfish in the wild. *FISH. RES. J. PHILIPP.* 6 (1): 1-10
- Bautista MN, de la Cruz MC. 1988. Linoleic (omega 6) and linolenic (omega 3) acids in the diet of fingerling milkfish (*Chanos chanos* Forsskal). *AQUACULTURE* 71 (4): 347-358
- Bautista MN, del Valle MJ, Orejana FM. 1991. Lipid and fatty acid composition of brackishwater and freshwater-reared milkfish (*Chanos chanos* Forsskal). *AQUACULTURE*. 96 (3-4): 241-248
- Benitez LV, Tiro LB. 1982. Studies on the digestive proteases of the milkfish *Chanos chanos*. *MAR. BIOL.* 71 (3): 309-315
- Borlongan IG. 1992. The essential fatty acid requirement of milkfish (*Chanos chanos* Forsskal). *FISH PHYSIOL. BIOCHEM.* 9 (5-6): 401-407
- Borlongan IG. 1992. Dietary requirement of milkfish (*Chanos chanos* Forsskal) juveniles for total aromatic amino acids. *AQUACULTURE* 102 (4): 309-317
- Borlongan IG. 1991. Arginine and threonine requirements of milkfish (*Chanos chanos* Forsskal) juveniles. *AQUACULTURE* 93 (4): 313-322
- Borlongan IG. 1990. Studies on the digestive lipases of milkfish, *Chanos chanos*. *AQUACULTURE* 89 (3-4): 315-325
- Buri P. 1981. Pigmentation pattern in the early developmental stages of milkfish (*Chanos chanos*): A key for larval identification. *FISH. RES. J. PHILIPP.* 6 (1): 51-55
- Buri P, Banada V, Triño A. 1981. Developmental and ecological stages in the life history of milkfish *Chanos chanos* (F.). *FISH. RES. J. PHILIPP.* 6 (2): 33-58
- Chaudhuri H, Juario JV. 1977. Use of hormones in breeding cultivated warm-water fishes with special reference to milkfish, *Chanos chanos* (Forsskal). *FISH. RES. J. PHILIPP.* 2 (2): 1-6
- Chaudhuri H, Juario J, Samson R, Tiro L. 1976. Notes on the external sex characters of *Chanos chanos* (Forsskal) spawners. *FISH. RES. J. PHILIPP.* 1 (2): 76-85

- Chaudhuri H, Juario JV, Primavera JH, Samson R, Mateo R. 1978. Observations on artificial fertilization of eggs and the embryonic and larval development of milkfish *Chanos chanos* (Forsskal). AQUACULTURE 13 (2): 95-113
- Chiu YN, Benitez LV. 1981. Studies on the carbohydrases in the digestive tract of the milkfish *Chanos chanos*. MAR. BIOL. 61 (2-3): 247-254
- Coloso RM, Tiro LB, Benitez LV. 1992. Requirement for tryptophan by milkfish (*Chanos chanos* Forsskal) juveniles. FISH PHYSIOL. BIOCHEM. 10 (1): 35-41
- de la Peña LD, Chiu YN, Gancharo F. 1987. Evaluation of various leguminous seeds as protein sources for milkfish, *Chanos chanos* Forsskal, juveniles. ASIAN FISH. SCI. 1 (1): 19-25
- Duray MN. 1995. The effect of tank color and rotifer density on rotifer ingestion, growth and survival of milkfish (*Chanos chanos*) larvae. PHILIPP. SCI. 32 18-26
- Duray M, Bagarinao T. 1984 Weaning of hatchery-bred milkfish larvae from live food to artificial diets. AQUACULTURE 41 (4): 325-332
- Emata AC, Marte CL. 1994. Natural spawning, egg and fry production of milkfish, *Chanos chanos* (Forsskal), broodstock reared in concrete tanks. J. APPL. ICTHYOL. 10 (1): 10-16
- Emata AC, Marte CL. 1993. Broodstock management and egg production of milkfish, *Chanos chanos* Forsskal. AQUACULT. FISH MANAGE. 24 (3) 381-388
- Emata AC, Marte CL. 1992. The use of a visual implant tag to monitor the reproductive performance of individual milkfish *Chanos chanos* Forsskal. J. APPL. ICTHYOL. 8 (1-4): 314-317
- Emata AC, Marte CL, Garcia LMB. 1992. Management of milkfish broodstock. Tigbauan, Iloilo, Philippines: SEAFDEC Aquaculture Department. 22 pp
- Estudillo CB, MN Duray, Marasigan ET. 1998. Growth and survival of milkfish (*Chanos chanos*), sea bass (*Lates calcarifer*), and rabbitfish (*Siganus guttatus*) larvae reared at the same density in different sized tanks. ISR. J. AQUACULT. - BAMIDGEH 50: 20-24
- Gapasin RSJ, Bombeo R, Lavens P, Sorgeloos P, Nelis H. 1998. Enrichment of live food with essential fatty acids and vitamin C: effects on milkfish (*Chanos chanos*) larval performance. AQUACULTURE 162 (3-4): 271-288
- Garcia LMB, Gapasin RSJ. 1988. An inexpensive tag for short-term studies in milkfish (*Chanos chanos* Forsskal) and in seabass (*Lates calcarifer* Bloch). J. APPL. ICTHYOL. 4 (3): 101-104
- Garcia LMB, Toledo JD 1988. Critical factors influencing survival and hatching of milkfish (*Chanos chanos* Forsskal) eggs during simulated transport. AQUACULTURE 72 (1-2) 85-93
- Hara S, Canto JT Jr., Almendras JME. 1982. A comparative study of various extenders for milkfish, *Chanos chanos* (Forsskal), sperm preservation. AQUACULTURE 28 (3-4): 39-346
- Hilomen-Garcia G.V. 1998. Sensitivity of fertilized milkfish (*Chanos chanos* Forsskal) eggs to mechanical shock and simulated transport. AQUACULTURE 159 (3-4): 239-247
- Hilomen-Garcia GV. 1997. Morphological abnormalities in hatchery-bred milkfish (*Chanos chanos* Forsskal) fry and juveniles. AQUACULTURE 152 (1-4): 155-166
- Juario JV, Storch V. 1984. Biological evaluation of phytoplankton (*Chlorella* sp., *Tetraselmis* sp. and *Isocrysis galbana*) as food for milkfish (*Chanos chanos*) fry. AQUACULTURE. 40 (3) : 193-198
- Juario JV, Quintio GF, Banno JE, Natividad M. 1980. Effects of exogenous hormone injections on milt consistency in newly caught, wild milkfish. KALIKASAN 9 (2-3): 321-326

- Ferraris RP, Almendras JM, Jazul AP. 1988. Changes in plasma osmolality and chloride concentration during abrupt transfer of milkfish (*Chanos chanos*) from seawater to different test salinities. *AQUACULTURE* 70 (1-2): 145-157
- Ferraris RP, Catacutan MR, Mabelin RL, Jazul AP. 1986. Digestability in milkfish, *Chanos chanos* (Forsskal): Effects of protein source, fish size and salinity. *AQUACULTURE* 59 (2): 93-105
- Kumagai S, Castillo NM. 1980. Changes in the length and weight of milkfish, *Chanos chanos*, larvae preserved in formalin. *FISH. RES. J. PHILIPP.* 5 (1): 17-23
- Lim C, Sukhawongs S, Pascual FP. 1979. A preliminary study on the protein requirements of *Chanos chanos* (Forsskal) fry in a controlled environment. *AQUACULTURE* 17 (3): 195-201
- Marte CL. 1998. An improved method for collecting naturally spawned milkfish eggs from floating cages. *AQUACULTURE* 71 (4): 387-392
- Marte CL, Lacanilao F. 1986. Spontaneous maturation and spawning of milkfish in floating net cages. *AQUACULTURE* 53 (2): 115-132
- Marte CL, Lam TJ. 1992. Hormonal changes accompanying sexual maturation in captive milkfish (*Chanos chanos* Forsskal). *FISH PHYSIOL. BIOCHEM.* 10 (4): 267-275
- Marte CL, Sherwood NM, Crim LW, Harvey B. 1987. Induced spawning of maturing milkfish (*Chanos chanos* Forsskal) with gonadotropin-releasing hormone (GnRH) analogues administered in various ways. *AQUACULTURE* 60 (3-4): 303-310
- Pantastico JB, Baldia JP, Reyes DM Jr. 1986. Feed preference of milkfish (*Chanos chanos* Forsskal) fry given different algal species as natural feed. *AQUACULTURE* 56 (3-4): 169-178
- Santiago CB, Banes-Aldaba M, Songalia ET. 1983. Effect of artificial diets on growth and survival of milkfish fry in water. *AQUACULTURE* 34 (3-4): 247-252
- Santiago CB, Pantastico JB, Baldia SF, Reyes OS. 1989. Milkfish (*Chanos chanos*) fingerling production in freshwater ponds with the use of natural and artificial feeds. *AQUACULTURE* 77 (4): 307-318
- Toledo JD, Gaitan AG. 1992. Egg cannibalism by milkfish (*Chanos chanos* Forsskal) spawners in circular floating net cages. *J. APPL. ICTHYOL.* 8 (1-4): 257-262
- Villaluz AC, Unggui A. 1983. Effects of temperatures on behavior, growth, development and survival in young milkfish, *Chanos chanos* (Forsskal). *AQUACULTURE* 35 (4): 321-330
- Villegas CT. 1990. The effects on growth and survival of feeding water fleas (*Moina macrocopa* Strauss) and rotifers (*Brachionus plicatilis*) to milkfish (*Chanos chanos* Forsskal) fry. *ISR. J. AQUACULT.-BAMIDGEH* 42 (1) 10-17
- Villegas CT, Lumasag GL. 1991. Biological evaluation of frozen zooplankton as food for milkfish (*Chanos chanos*) fry. *J. APPL. ICTHYOL* 7 (2): 65-71
- Villegas CT, Bombeo T. 1982. Effects of increased stocking density and supplemental feeding on the production of milkfish fingerlings. *FISH. RES. J. PHILIPP.* 7 (2) 21-27
- Villegas CT, Millamena O, Escritor F. 1990. Food value of *Brachionus plicatilis* fed three selected algal species as live food for milkfish, *Chanos chanos* Forsskal, fry production. *AQUACULT. FISH. MANAGE.* 21 (2): 213-219

FOR COPIES OF THESE REPORTS

**E-mail: library@aqd.seafdec.org.ph
aqdlib@i-iloilo.com.ph
Fax: (63 33) 335 1008, 336 2891**