

1978

The use of decapsulated brine shrimp eggs as food for shrimp larvae

Laviña, E.

Aquaculture Department, Southeast Asian Fisheries Development Center

Laviña, E., & Figueroa, R. (1978). The use of decapsulated brine shrimp eggs as food for shrimp larvae. SEAFDEC Aquaculture Department Quarterly Research Report, 2(4), 11–14.

<http://hdl.handle.net/10862/2334>

Downloaded from <http://repository.seafdec.org.ph>, SEAFDEC/AQD's Institutional Repository

The use of decapsulated brine shrimp eggs as food for shrimp larvae

E. Laviña and R. Figueroa

In most shrimp hatcheries throughout the world, freshly hatched nauplii of the brine shrimp *Artemia salina* are being fed to shrimp larvae, fish and other animal larvae.

Sorgeloos et. al (1977), have improved on the utilization of brine shrimp in aquaculture by modifying the techniques of Nakanishi et. al and Morris and Afzelius by removing the outer shell layer of *Artemia* cysts. This decapsulation technique removes the tedious and cumbersome process of separating freshly hatched nauplii from the empty shells and other debris, disinfects the cysts from pathogens and makes possible the direct feeding of decapsulated cysts to fish and crustacean larvae. This study was made to test the acceptability of brine shrimp decapsulated eggs to the larvae of some commercially important shrimp species. Direct feeding of decapsulated eggs would eliminate the process of incubating the eggs thereby cutting on labor and energy costs.

Decapsulated cysts of *Artemia salina* were experimentally fed to the larvae of *Penaeus monodon*, *Metapenaeus ensis*, *M. endeavouri* and *Macrobrachium rosenbergii* and were found to be comparable to freshly hatched brine shrimp nauplii.

Results of the experiment are presented in Table 1. A single classification ANOVA for the results of each experiment show significance at $P < 0.05$ for *P. monodon*, *P. indicus*, *M. ensis* and *M. rosenbergii* and a high significance at $P < 0.01$ for *P. indicus*, *M. ensis* and *M. rosenbergii*. Survival of the experimental and control groups in *M. endeavouri* is not significant.

Table 2 shows the mean number of cyst/nauplii consumed per individual per treatment in each experiment. In all experiments, the Student-Newman-Keuls test to compare the mean of the samples show significance at $P < 0.05$ between the control and experimental groups. No significant difference was found between the mean of the groups fed with decapsulated cysts and of those fed with freshly hatched nauplii. There was no significant difference in the number of decapsulated cysts or nauplii consumed by the shrimp larvae. The consumption rate of each species however varied.

Direct feeding of decapsulated cysts to shrimp and fish larvae eliminates the process of incubation and hatching of brine shrimp cysts thereby reducing labor costs and simplifying the preparations for larval feeding. Furthermore, decapsulated cysts can be conveniently stored at low temperatures. Decapsulated cysts in diapause which might possibly fail to hatch after incubation become available as food to the shrimp and fish larvae thus reducing losses in feed.

It must be noted that the decapsulated brine shrimp cysts lose some buoyancy due to loss of their outer chorion during decapsulation and tend to sink in seawater. Continuous suspension of the decapsulated cysts when provided with aeration from the bottom or with airwater lift pipes will avoid such problem.

Table 1. Survival of the larvae of *Penaeus monodon*, *P. indicus*, *Metapenaeus ensis*, *M. endeavori*, and *Macrobrachium rosenbergii* fed daily with 2,000 Jecapsulated cysts (Group B) or 2,000 freshly hatched brine shrimp (Group C). The controls (Group A) and all the other group samples were each given about 15×10^6 *Chaetoceros calcitrans* cells/days

Expt. no	Species	Dates	Initial no. of test ind.	Replicates	Treatments			Calc. Fs value
					A	B	C	
I	<i>P. monodon</i>	8/10 to 8/13/77	30	1	15	27	28	8.61 *
				2	22	28	23	
				3	20	28	29	
				Ave.	19	27.67	26.67	
II	<i>P. indicus</i>	7/13 to 7/18/77	20	1	0	17	15	29.10 **
				2	4	14	10	
				3	0	13	13	
				Ave.	1.33	14.67	12.67	
III	<i>M. ensis</i>	7/23 to 7/30/77	30	1	4	24	27	71.896 **
				2	5	29	21	
				3	2	25	26	
				Ave.	3.67	26	24.67	
IV	<i>M. endeavori</i>	7/25 to 8/1 /77	30	1	24	25	21	0.37
				2	20	12	17	
				3	21	20	20	
				Ave.	21.67	19	19.3	
	<i>M. rosenbergii</i>	7/25 to 8/1 /77	30	1	8	26	28	126.51 **
				2	11	29	28	
				3	12	27	27	

* Significant at $P < 0.05$

** Highly significant at $P < 0.01$

Tabulated F value at 5% confidence limit is 5.14

Tabulated F value at 1% confidence limit is 10.92

Table 2. Mean number of cysts/nauplii consumed per individual per treatment per day

Expt. no.	Species	Dates	Replicates	Treatments		Calc. F_s values
				B	C	
I	<i>P. monodon</i>	8/10 to 8/13/77	1	62.172	68.24	.779
			2	72.05	83.66	
			3	69.03	67.31	
			Ave.	70.084	73.07	
II	<i>P. indicus</i>	7/13 to 7/18/77	1	115.1412	109.14	.169
			2	117.96	122.29	
			3	132.39	125.115	
			Ave.	121.83	118.85	
III	<i>M. ensis</i>	7/23 to 7/30/77	1	73.46	70.93	2.396
			2	66.49	80.34	
			3	62.93	71.68	
			Ave.	67.63	74.32	
IV	<i>M. endeavori</i>	7/25 to 8/1/77	1	73.43	78.30	.459
			2	107.5	88.37	
			3	90.9	84.30	
			Ave.	90.61	83.66	
V	<i>M. rosenbergii</i>	7/25 to 8/1 /77	1	65.99	64.43	.260
			2	65.77	64.31	
			3	66.12	67.72	
			Ave.	65.96	66.49	

All F_s values not significant at $P < 0.05$
 Tabulated F at 5% confidence limit is 7.71

LITERATURE CITED

- Bardach, J. E., J. H. Ryther, and W. O. McLaren. 1972. Aquaculture: farming and husbandry of freshwater and marine organism. Wiley Interscience, New York, 868 pp.
- Platon, R. 1977. Experimental Hatchery Project. Third Quarterly Report, SEAFDEC, Aquaculture Department.
- Sorgeloos, P. 1976. The brine shrimp, *Artemia salina* L.: A bottleneck in Mariculture, FAO Techn. Conf. on Aquaculture, Kyoto (Japan), Experience paper 77:12 p.
- Sorgeloos, P., E. Bossuyt, E. Lavina, M. Baeza-Mesa, and G. Persoone. 1977. Decapsulation of *Artemia* cysts: a simple technique for the improvement of the use of brine shrimp in aquaculture. *Aquaculture*, 12:311-316.
- Sorgeloos, P. and G. Persoone. 1975. Technological improvements for the cultivation of invertebrates as food for fishes and crustaceans. II. Hatching and culturing of the brine shrimp *Artemia salina* L. *Aquaculture*, 6:303-317.
- Santos, C. de los, Sorgeloos, P., Lavina, E., and A. Bernardino. 1979. Successful inoculation of *Artemia* and cysts production in man-made salterns. Int'l Conference *Artemia salina*, Corpus Christi, Texas, U.S.A. 20-23 August 1979.
- Villaluz, D. K., A. Villaluz, B. Ladrera, M. Sheik and A. Gonzaga. 1972. Production, larval development and cultivation of sugpo (*Penaeus monodon* Fabricius). *Phil. J. of Science*. 98 (3-4): 205-233.